

SCIPP 93/27
February 1, 2008

INVESTIGATION OF DISCRETE GAUGE ANOMALIES IN STRING THEORY[★]

D. A. MACINTIRE[†]

*Santa Cruz Institute for Particle Physics
University of California, Santa Cruz, CA 95064*

Abstract

In string theory, there are no continuous global symmetries. Discrete symmetries frequently appear, and these can often be understood as unbroken subgroups of larger, spontaneously broken gauge symmetries (discrete gauge symmetries). In cases which have been studied previously, anomalies in these symmetries could always be cancelled by a Green-Schwarz mechanism. In the present work, we describe results of an investigation of a large number of $\mathbb{Z}_3$ orbifold models with two and three Wilson lines. We find that the discrete gauge anomalies can always be cancelled.

[★] Work supported in part by the U.S. Department of Energy.

[†] Address after September 1, 1993: Department of Physics, University of Colorado, Boulder, CO 80309.

In the absence of a definitive understanding of the vacuum structure of string theory, it is of interest to enumerate those properties of the theory that may be considered generic or universal. One well known example^[1] is that all continuous symmetries in string theory are gauge symmetries. This remarkably powerful conclusion is reached by noting that for every current there is a corresponding gauge boson vertex operator. Whether or not this conclusion can be extended to *discrete* symmetries is of great interest. In field theory, gauged discrete symmetries,^[2] by definition, arise through the spontaneous breakdown of continuous local symmetries. It is clear then that any such discrete symmetry is anomaly-free, this being inherited from the underlying symmetry.

At low energies, this anomaly freedom leads to a useful constraint obtained by requiring that the 't Hooft effective fermion vertex be invariant under the discrete symmetry. Otherwise, instanton effects would lead to explicit violations of the symmetry, and it could not be gauged. In string theory, previous studies suggest that a slightly more general result holds.^[3] In this case the instanton vertex is accompanied by an exponential of the model-independent axion field (the partner of the dilaton, whose vacuum expectation value sets the value of the gauge coupling). A discrete symmetry can be gauged only if the transformation of the fermionic part of the instanton vertex can be cancelled by assigning a shift to the model-independent axion. This is just the four-dimensional analogue of the Green-Schwarz mechanism applied to discrete symmetries. In a large number of Calabi-Yau compactifications (with or without Wilson lines), it has been shown that such an assignment is possible.^[4] We should note, however, that there is no deep understanding of how this transformation law arises. (Nor can one be certain that it *does* arise.) This is to be contrasted with the Green-Schwarz terms for continuous symmetries, where one can check that the appropriate couplings (to gauge bosons) appear in perturbation theory.

This cancellation is non-trivial in the case in which there is more than one non-Abelian gauge group. The discrete anomalies of each gauge group must match in order that they all be cancelled by the transformation of the one model-independent

axion. If all discrete symmetries are gauge symmetries in string theory, then they must pass this rather stringent test.

In this paper we report the results of an investigation of discrete gauge anomalies in a large number of models orbifold models with Wilson lines. We find that in all models investigated, the discrete gauge anomalies can be cancelled through a transformation of the axion.

1. $\mathbb{Z}_3$ Orbifolds with Wilson Lines

The $\mathbb{Z}_3$ orbifold compactification of the heterotic string is simple and well known.^[5] We begin by reviewing the relevant formalism.

The $\mathbb{Z}_3$ orbifold with Wilson lines is obtained by modding $\mathbb{R}^6$ by the space group S , consisting of rotations in three two-dimensional planes and translations:

$$S : Z \rightarrow \Theta Z + v. \quad (1.1)$$

Associated with this is a transformation of the $E_8 \times E_8$ lattice

$$\begin{aligned} (\Theta, 0) &\rightarrow (\mathbb{1}, V^I) \\ (\mathbb{1}, e_m^i) &\rightarrow (\mathbb{1}, a_m^I) \end{aligned} \quad (1.2)$$

and a transformation of the bosonized fermion lattice:

$$\begin{aligned} (\Theta, 0) &\rightarrow (\mathbb{1}, v_R) \\ (\mathbb{1}, e_m^i) &\rightarrow (\mathbb{1}, 0). \end{aligned} \quad (1.3)$$

These models are completely specified by the shift vector $\mathbf{V}$ and Wilson lines $a_m^I, m = 1, 3, 5$. These vectors must satisfy the modular invariance constraints

$$3 \left(\mathbf{V} + \sum_m n^m \mathbf{a}_m \right)^2 = 0 \bmod 2, \quad n^m = 0, \pm 1 \quad (1.4)$$

and the requirements $3\mathbf{V}, 3\mathbf{a}_m \in E_8 \times E_8$. The gauge bosons for these models

correspond to vectors $\mathbf{P} \in E_8 \times E_8$ satisfying

$$\mathbf{P}^2 = 2, \quad \mathbf{P} \cdot \mathbf{V} \in \mathbb{Z}, \quad \mathbf{P} \cdot \mathbf{a}_m \in \mathbb{Z}, \quad \forall m. \quad (1.5)$$

Massless matter states from the untwisted sector are given by

$$\mathbf{P}^2 = 2, \quad \mathbf{P} \cdot \mathbf{V} = \frac{1}{3} \bmod 1, \quad \mathbf{P} \cdot \mathbf{a}_m \in \mathbb{Z}, \quad \forall m. \quad (1.6)$$

Massless states from the twisted sectors are given by

$$\left(\mathbf{P} + \mathbf{V} + \sum_m n^m \mathbf{a}_m \right)^2 = \frac{4}{3} \quad (\text{multiplicity } 1) \quad (1.7)$$

and

$$\left(\mathbf{P} + \mathbf{V} + \sum_m n^m \mathbf{a}_m \right)^2 = \frac{2}{3} \quad (\text{oscillator states, multiplicity } 3). \quad (1.8)$$

With three inequivalent non-zero Wilson lines there are twenty-seven distinct twisted sectors.

Whereas the modding Θ is chosen to be the diagonal $\mathbb{Z}_3$ given by $\Theta_j^i = \delta_j^i \exp(2\pi i \eta^j / 3)$ with $\eta = (1, 1, -2)$, we will look for anomalies in the discrete $\mathbb{Z}_3$ rotation of *one* of the three $SU(3)$ lattices, referred to here as G_1 . Before modding, it is clear that this is just a general coordinate transformation in the internal space. That it remains a symmetry of the interacting theory may be verified by checking that it commutes with the BRST operator. To determine the action of this discrete rotation on the fermions of the model, it is simplest to bosonize the right-moving fermions. In the light-cone gauge, the quantized momenta of bosons describing the (GSO projected) Ramond sector belong to the (s) representation of $SO(8)$. The positive helicity momenta are therefore of the form

$$\mathbf{k} = \left(n_1 + \frac{1}{2}, n_2 + \frac{1}{2}, n_3 + \frac{1}{2}; +\frac{1}{2} \right), \quad \sum_i n_i = 0 \bmod 2, \quad n_i \in \mathbb{Z} \quad (1.9)$$

where the last component of $\mathbf{k}$ is the spacetime helicity.

A G_1 -rotation of $2\pi/3$ of the first $SU(3)$ lattice corresponds to a shift of the bosonized field:

$$\phi \rightarrow \phi + 2\pi \mathbf{v}_1 \quad \mathbf{v}_1 = \left(\frac{1}{3}, 0, 0; 0\right). \quad (1.10)$$

The right-moving component of a state will therefore acquire a phase under the discrete G_1 -rotation given by $e^{2\pi i \mathbf{k} \cdot \mathbf{v}_1}$. We see also from this consideration that G_1 is an R-symmetry; the gauginos (and the supercharges), with $\mathbf{k} = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}; \frac{1}{2})$, transform under G_1 by a phase $e^{i\pi/3}$.

In the twisted sectors, the momenta (1.9) are shifted according to

$$\mathbf{k} \rightarrow \mathbf{k} + \mathbf{v}_R \quad \mathbf{v}_R = \left(\frac{1}{3}, \frac{1}{3}, -\frac{2}{3}; 0\right). \quad (1.11)$$

Massless twisted (matter) sector states must have a right-moving component with momentum given by[★]

$$\mathbf{k} + \mathbf{v}_R = \left(\frac{1}{6}, \frac{1}{6}, \frac{1}{6}; -\frac{1}{2}\right). \quad (1.12)$$

These states acquire a phase under the discrete rotation given by $e^{\{2\pi i (\mathbf{k} + \mathbf{v}_R) \cdot \mathbf{v}_1\}}$.

G_1 will also have an effect on the left-moving component of the massless oscillator states of the form $|\mathbf{k}\rangle_R \tilde{\alpha}_{-1/3}^i |\mathbf{P}\rangle_L$, $i = 1, 2, 3$. Because $\tilde{\alpha}^i$ (in particular, $\tilde{\alpha}^1$) transforms like Z^i under the discrete rotation, there is an additional phase of $e^{2\pi i/3}$ that must be taken into account for states of this type.

★ We are indebted to Luis Ibanez for pointing out that one must be careful with this helicity assignment for the twisted sector.

2. Models

The 't Hooft vertex for a given model and non-Abelian gauge group is a product of the form

$$e^{ia/f_a} \prod_{massless} \psi\psi \cdots \psi$$

with one field ψ for each zero mode of a given multiplet. There is a single zero mode for each doublet of $SU(2)$ or triplet of $SU(3)$, and four or six zero modes for the $SU(2)$ or $SU(3)$ gauginos respectively. Defining $\alpha = e^{2\pi i/3}$ and using the above relations, we find that for a model with N_{UT} untwisted matter doublets or triplets, and N_{TW} matter doublets or triplets, the transformation of the 't Hooft vertex under the discrete transformation is given by

$$e^{ia/f_a} \prod_{massless} \psi\psi \cdots \psi \rightarrow \alpha^{-N_{UT}/6} \alpha^{N_{TW}/6} \alpha^{C_G} e^{ia/f_a} \prod_{massless} \psi\psi \cdots \psi \quad (2.1)$$

with C_G the quadratic Casimir. If the phase given in Eq. (2.1) is non-zero, the discrete symmetry is anomalous. However, note the presence of the model-independent axion in the 't Hooft vertex; its form is universal for each non-Abelian group. The discrete anomaly may be cancelled by assigning a G_1 -transformation to the axion if and only if the transformation (2.1) is the *same* for each non-Abelian group. In earlier studies, it was shown that many models do satisfy this condition. For example, if one takes the Calabi-Yau manifold Y_5^4 , with Wilson lines, one finds that this condition is satisfied for any choice of Wilson line, for either the $E_8 \times E_8$ or $O(32)$ theories.

An example of a particular model with cancellable discrete gauge anomalies is generated by the vectors:

$$\begin{aligned} \mathbf{V} &= \left(-\frac{2}{3}, 0, -\frac{2}{3}, 0, 0, -\frac{2}{3}, 0, 0\right) \left(0, 0, -\frac{2}{3}, \frac{2}{3}, \frac{2}{3}, 0, 0, 0\right) \\ \mathbf{a}_1 &= \left(0, 0, 0, \frac{2}{3}, 0, 0, 0, 0\right) \left(0, \frac{1}{3}, 0, 0, 0, 0, -\frac{1}{3}, 0\right) \\ \mathbf{a}_3 &= \left(0, 0, 0, -\frac{1}{3}, \frac{2}{3}, \frac{1}{3}, \frac{2}{3}, -\frac{2}{3}\right) \left(-\frac{1}{3}, \frac{2}{3}, \frac{2}{3}, 0, 0, 0, 0, -\frac{1}{3}\right) \\ \mathbf{a}_5 &= \left(0, 0, 0, 0, 0, \frac{2}{3}, 0, 0\right) \left(0, 0, -\frac{1}{3}, 0, \frac{1}{3}, 0, 0, 0\right). \end{aligned} \quad (2.2)$$

This model has the gauge group $SU(3) \otimes SU(2) \otimes SU(2)$ with $SU(2)$ charge operators given by:

$$\begin{aligned} I_3 &= \frac{1}{2}(1, 0, -1, 0, 0, 0, 0, 0)(0, 0, 0, 0, 0, 0, 0, 0) \\ I'_3 &= \frac{1}{2}(0, 0, 0, 0, 0, 0, 0, 0)(1, 0, 0, 0, 0, 0, 0, -1) \end{aligned}$$

The full spectrum of this model is

$$(\mathbf{3}, \mathbf{1}, \mathbf{1}) + (\bar{\mathbf{3}}, \mathbf{1}, \mathbf{1}) + 20 [(\mathbf{1}, \mathbf{2}, \mathbf{1}) + (\mathbf{1}, \mathbf{1}, \mathbf{2})] + 205(\mathbf{1}, \mathbf{1}, \mathbf{1}). \quad (2.3)$$

Six of the twenty $SU(2)$ doublets are from the untwisted sector, and the remaining doublets are from the twisted sector, for both $SU(2)$'s. For the $SU(3)$, both the $\mathbf{3}$ and $\bar{\mathbf{3}}$ are twisted matter. The discrete transformation for the fermionic part of the 't Hooft action for each of the gauge groups is $\alpha^{1/3}$.

On the order of 10^5 randomly generated $\mathbb{Z}_3$ orbifold models were investigated. The search included models with two or three non-zero Wilson lines. We found that in all cases the discrete anomalies can be cancelled through a transformation of the model independent axion.

3. Conclusions

Previously, only a limited number of string theory models have been checked for the cancellation of discrete gauge anomalies. In this paper we report on an investigation of a huge number of models, and have found that this cancellation always occurs. While searches to date have involved only particular types of models, and the search should be expanded to other types of string theory models, we suspect that there exists some general connection between the requirement of modular invariance and the cancellation of discrete gauge anomalies.

REFERENCES

1. T. Banks and L. Dixon, *Nucl. Phys.* **B307** (1988) 93.
2. L. Krauss and F. Wilczek, *Phys. Rev. Lett.* **62** (1989) 1221; L.E. Ibáñez and G.G. Ross, *Phys. Lett.* **260B** (1991) 291; *Nucl. Phys.* **B368** (1992) 3; J. Preskill, S. Trivedi, F. Wilczek and M. Wise, *Nucl. Phys.* **B363** (1991) 207.
3. T. Banks and M. Dine, *Phys. Rev.* **D45** (1992) 424.
4. M. Dine, R. Leigh, D.A. MacIntire (unpublished).
5. L. Dixon, J.A. Harvey, C. Vafa and E. Witten, *Nucl. Phys.* **B261** (1985) 678; *Nucl. Phys.* **B274** (1986) 285; L.E. Ibáñez, H.P. Nilles, F. Quevedo, *Phys. Lett.* **187B** (1987) 25.